

From the Cosmological Microwave Background to Large-Scale Structure

Joseph Silk and Eric Gawiser

*Departments of Astronomy and Physics, and Center for Particle Astrophysics,
University of California, Berkeley, CA 94720*

The shape of the primordial fluctuation spectrum is probed by cosmic microwave background fluctuations which measure density fluctuations at $z \sim 1000$ on scales of hundreds of Mpc and from galaxy redshift surveys, which measure structure at low redshift out to several hundred Mpc. The currently acceptable library of cosmological models is inadequate to account for the current data, and more exotic models must be sought. New data sets such as SDSS and 2DF are urgently needed to verify whether the shape discrepancies in $P(k)$ will persist.

1 Introduction

Our understanding of primordial fluctuations in the early universe was revolutionized first with inflation and then by the actual detection of temperature fluctuations in the cosmic microwave background (CMB). Inflation gave the spectrum of fluctuations, but not the normalization. The COBE-DMR experiment measured the amplitude of temperature fluctuations on an angular scale that was acausal at last scattering, and hence directly probed inflationary fluctuations, and in particular the fluctuation strength. More than twenty subsequent experiments have plugged the causal gap, measuring fluctuations over angular scales less than or of order the acoustic peak at $\lambda = 220 \Omega^{1/2}$ that corresponds to the maximum sound horizon in the early universe.

The shape of the fluctuation spectrum is now being probed. The CMB fluctuations measure density fluctuations at $z \sim 1000$ on scales of hundreds of Mpc. At large redshifts one degree subtends a comoving scale of 100 Mpc. A complementary measure arises from galaxy redshift surveys. These measure variations in the luminous matter density out to ~ 300 Mpc at the present epoch. One can combine, after choosing a model, the CMB and large-scale structure (LSS) measures of the fluctuation spectrum. Here we will describe the current status of our understanding of the shape of the primordial fluctuation spectrum.

It is customary to use a two-parameter fit to the LSS power: σ_8 , the normalization at $8 h^{-1}$ Mpc, and Γ , measure of shape relative to CDM and nearly equal to Ωh for CDM. Given the several data sets, each with a number of independent data points, this may be an unnecessarily restrictive approach. Of course any data set is imperfect, with possible systematic errors, and the

data sets have different selection biases. However, provided the bias is scale-independent, one can renormalize the data sets and examine detailed shape constraints. One has to decide whether to compare a nonlinear power spectrum with the data or whether to correct the data for nonlinearity and compare the data with linear theory. We will employ the latter approach here.

2 CMB: Status of the Theory

Inflation-generated curvature fluctuations provide the paradigm for interpreting the cosmic microwave background temperature anisotropies. There are three components to $\delta T/T$, schematically summarized as

$$\frac{\delta T}{T} = |\delta\phi + \delta\rho_\gamma + \delta v|.$$

These are the gravitational potential, intrinsic and Doppler contributions from the last scattering surface. The combined effect of the first two terms results in the Sachs-Wolfe effect $\delta T/T = \frac{1}{3}|\delta\phi|$ which represents the only superhorizon contributions to $\delta T/T$. Inflationary initial conditions then require that the Fourier component $|\delta T/T|_k \propto |\cos(kv_s t_{ls})|$ at the last scattering epoch t_{ls} , where k is the wavenumber and v_s is the sound speed, must be constant as $k \rightarrow 0$. Inflation of course specifies the phases of density fluctuations that decompose to sound waves of wavelength less than that of the maximum sound horizon, $v_s t_{ls}$. Longer wavelengths correspond to power-law growing modes at horizon crossing. The wave just entering the horizon at last scattering has a peak at wavenumber $n\pi/v_s t_{ls}$, $n = 1$, and a succession of waves crest at $n = 2, 3, \dots$ before damping sets in as the photon mean free path increases relative to the wavelength. The first acoustic peak projects to $\delta T/T$ on angular scales $\sim \Omega^{1/2}$ degree, and is a robust measure of the curvature of the universe. Doppler peaks are 90° out of phase and of lower amplitude, so they fill in the troughs of the acoustic oscillations as measured by the radiation power spectrum. Peak heights are determined in large part by choice of Ω_B and Ω_Λ . An increase in Ω_B enhances the wave compression and reduces the rarefaction phases. An increase in Ω_Λ enhances the ratio of radiation to matter in a flat model, and thereby boosts the peak potential decay and the low ℓ power via the early integrated Sachs-Wolfe effect. Of course increasing the spectral index n also raises the peak height. Peak heights are lowered by reionization and secondary scattering. Not all of these degeneracies are removed by examining the higher peaks. For example, combination of Ω_B and Ω_Λ at specified Ω is nearly degenerate in peak height and peak location since the angular size-redshift relation depends only on Ω .

Lensing by nonlinear and quasilinear foreground structure redistributes the peak power towards very high ℓ in a way that breaks the degeneracies in the CMB. One can search for this effect with interferometer experiments¹ at $\ell \gtrsim 1000$ or else by correlating the CMB fluctuations with large-scale power from either large redshift surveys such as the SDSS² or via weak lensing distortions of the CMB³ or LSS⁴.

3 Reconstruction of the Primordial Power Spectrum from the CMB

The inflationary, approximately scale-free, power spectrum $P(k) = Ak^n$, $n \approx 1$, is modified by the transition from radiation to matter domination, since in matter-dominated epochs different growth occurs on subhorizon scales: in the radiation era, there is no subhorizon growth of fluctuations. This modifies $P(k)$ to $P(k) \propto k^{-3}$ on larger scales. The transition occurs at the horizon during matter-radiation equality, namely $12(\Omega h^2)^{-1}$ Mpc. The Boltzmann equation can be solved for temperature fluctuations, mode by mode, and the solutions for $\delta T/T$ are scaled to agree with the quoted errors for each CMB experiment. For each specified cosmological model, we then infer the power spectrum amplitude over scales that correspond to the deprojection on the sky of the experimental window function⁶. We confirm that standard CDM ($\Omega_{CDM} = 1$, $h = 0.5$) fits the CMB data rather poorly, with best fit renormalization that corresponds to a high value of σ_8 . The Λ CDM model ($\Omega = 1$, $\Omega_m = 0.4$, $h = 0.6$) gives a reasonably good fit with an acceptable value of σ_8 . Much stronger constraints however come when these fits are combined with LSS data.

4 Reconstruction of $P(k)$ from Large-Scale Structure Data

There are several large-scale structure data sets that one may use to reconstruct $P(k)$. Galaxy redshift surveys include the Las Campanas Survey of 25,000 galaxies⁷, the PSCz survey of 1,500 galaxies⁸, and the SSRS2/CfA2 survey of 7,000 galaxies⁹. There is also the APM cluster survey which probes to $\sim 300 h^{-1}$ Mpc¹⁰ and the real space inversion of the 2D APM galaxy survey¹¹. The local mass function of clusters¹² has been used to measure $\sigma_8 \Omega^{0.6}$, and the high redshift cluster abundance¹³ has been used to break the degeneracy between σ_8 and Ω . Peculiar velocities and large scale bulk flows also yield $\sigma_8 \Omega^{0.6}$ in a completely bias-independent approach, although systematic uncertainties remain large¹⁴.

The redshift space surveys can be corrected in a straightforward way, for peculiar velocities on small scales and bulk flows on large scales, to derive the

real space $P(k)$ by assuming a cosmological model¹⁵. Correction of data in the nonlinear regime is best done by numerical simulation, but can be performed using an empirical formulation calibrated to numerical simulations based on a smooth interpolation from spherical collapse by a factor of 2 in radius on cluster scales¹⁶. Renormalization of the various measures of $P(k)$ is effected by assuming that all measurements are subject to a scale-independent bias, allowed to be independent for each probe of $P(k)$.

5 Confrontation of $P(k)$ with CMB and LSS

Model fitting to LSS alone results in the following conclusions. Of course the standard COBE-normalized CDM model fails completely. Without a large scale-dependent bias factor on 10 – 100 Mpc scales, peculiar velocities and the galaxy cluster abundance are greatly overpredicted. Low density models circumvent these problems. The cluster abundance, evolution and baryon fraction are all in satisfactory agreement with observations¹³.

However the combined CMB/LSS fits to $P(k)$ lead⁶ to a surprising conclusion. The surprise is that almost all models, while occasionally faring better than Λ CDM, still provide unacceptable fits to all of the data. Consider for example Λ CDM, currently favored by the SN Ia Hubble diagram. The reduced χ^2 is 2.1 for 70 degrees of freedom. Data set by data set, one still has a problem. For example the values of $\chi^2(\text{d.o.f.})$ are APM clusters: 25(8); LCRS 17(5); APM 44(9); IRAS 16(9). The acceptable data sets are CfA for 2 d.o.f., cluster abundances/peculiar velocities for 3 d.o.f. and CMB for 34 d.o.f.

6 Neutrinos and LSS

The only mildly acceptable model (reduced $\chi^2 = 1.2$) is CHDM, hot and cold dark matter with $\Omega = 1$. This model overpredicts current cluster abundances and underpredicts the small number of high redshift, luminous x-ray clusters (2 at $z = 0.5$, 1 at $z = 0.8$). However the cluster evolution constraint is disputed¹⁷, and the local normalization is not necessarily robust. The cluster baryon fraction provides an independent and powerful constraint that favors $\Omega_m \approx 0.3$. Of course this rests on the reasonably plausible assumption that clusters provide a fair sample of the baryon fraction of the universe. This need not necessarily be true if gas has had a complex history prior to cluster formation: *e.g.* the gas may have been preheated as is suggested by recent considerations of the entropy of intracluster gas¹⁸. This would reduce the baryon fraction, but one can equally well imagine scenarios for cluster formation in dense sheets or filaments where the baryon fraction was already enhanced.

Consider the model preferred by the combination of SNIa and cluster constraints, namely Λ CDM. Figure 1 shows the Λ CDM power spectrum compared with observations of Large-Scale Structure and CMB anisotropy. One can pose the following question: does adding a hot component improve the marginally acceptable LSS fit? We find that as an admixture of HDM is added to the dominant CDM, the combined fit to CMB and LSS deteriorates (Figures 2,3). The reason is that low density CDM models have a $P(k)$ peak that is longward of the apparent peak in the APM data. Adding HDM only exacerbates the mismatch.

Neutrino masses imprint a distinct signature on $P(k)$. This will eventually be a measurable probe of the neutrino mass, from LSS as well as from CMB. Indeed the LSS probe may potentially be more powerful¹⁹. There is more dynamical range available in probing $P(k)$ with LSS on the neutrino free streaming scale, where the primary signature should be present. Even present data is sensitive to a neutrino mass of around an eV: for example we find that the fit changes significantly between 0.1 and 1 eV. If Λ CDM is in fact the right model, our analysis yields an indirect upper limit on the mass of the most massive neutrino species of $m_\nu \leq 2\text{eV}$. While there are considerable systematic uncertainties in this approach, it is promising as a complement to the direct evidence for mass difference between neutrino species from SuperKamiokande²⁰ and the solar neutrino problem²¹, and is already beginning to conflict with results from LSND²² that require a large mass difference.

One class of exotic models is the following. Take a model that fits all constraints except for the shape. The best contender for such a model is Λ CDM. Inspection of the LSS constraints reveals that there is a deficiency of large-scale power near 100 Mpc. One can add an ad hoc feature on this scale from considering inflationary models with multiple scalar fields (see Figure 4). This could be generated for example by⁵ incomplete coagulation of bubbles of new phase in a universe that already has been homogenized by a previous episode of inflation. One can tune the bubble size distribution to be sharply peaked at any preferred scale. This results in nongaussian features and excess power where needed. The non-gaussianity provides a distinguishing characteristic.

Other suggestions that fit both CMB and LSS data appeal to an inflationary relic of excess power from broken scale invariance, arising from double inflation in a Λ CDM model, which results in a gaussian feature that is essentially a step in $P(k)$ at the desired wavenumber²³. This improves the fit in much the same way as adding a hot component to CDM improves the empirical fit. While such ad hoc fits may seem unattractive, one could argue that other aspects of cosmological model building are equally ad hoc, such as postulating a universe in which Λ is only becoming dynamically important at the present

epoch. Clearly one has to accommodate such arguments in order to fit the data, if the data is indeed accepted at face value. Moreover there are positive side effects that arise from the tuned void approach. The bubble-driven shells provide a source of overdensities on large scales. Rare shell interactions could produce nongaussian massive galaxies or clusters at low or even high redshift: above a critical surface density threshold gas cooling would help concentrate gas and aid collapse. If massive galaxies were discovered at say $z > 5$ or a massive galaxy cluster at $z > 2$ this would be another indication that the current library of cosmological models is inadequate. New data sets such as SDSS and 2DF are urgently needed to verify whether the shape discrepancies in $P(k)$ will persist.

7 Summary

If the data are accepted as mostly being free of systematics and ad hoc additions to the primordial power spectrum are avoided, there is no acceptable model for large-scale structure. No one LSS data set can be blamed. Perhaps it is best to wait for improved data. The Sloan and 2DF surveys are already acquiring galaxy redshifts. However another philosophy is to search for more exotic models. Consider for example the primordial isocurvature mode. This has the advantage of forming primordial black holes of stellar mass, since normalization to large-scale structure and the present spectrum over $10 - 50$ Mpc requires a spectral index that generates nonlinear fluctuations at roughly the epoch of the quark-hadron phase transition, when the horizon contained approximately one solar mass²⁴ (hence the primordial black holes may be the possibly observed MACHOs). The goodness of fit of this model to the combined CMB/LSS data is similar to that of the Λ CDM model. One cannot distinguish with current data between an exotic isocurvature model and Λ CDM, although neither model is satisfactory. To improve on this, clearly something even more exotic is required. It may be that independent observations will force us in this direction.

Acknowledgements

We gratefully acknowledge support by NSF and NASA.

References

1. B. Metcalf and J. Silk, ApJ 492L1 (1998)

2. Y. Wang, D. N. Spergel and M. A. Strauss, preprint astro-ph/9802231, ApJ, in press (1999)
3. M. Zaldarriaga and U. Seljak, preprint astro-ph/9810257 (1998)
4. W. Hu and M. Tegmark, preprint astro-ph/9811168 (1998)
5. L. Amendola, C. Baccigalupi, R. Konoplich, F. Occhionero and S. Rubin, Phys. Rev. D 54 7199 (1996)
6. E. Gawiser and J. Silk, Science 280 1405 (1998)
7. H. Lin et al., ApJ 471 617 (1996)
8. W. Saunders et al., Proc. of *XVIIth Rencontres de Moriond: Extragalactic Astronomy in the Infrared*, ed. G. A. Mamon, T. X. Thuan, J. T. T. Van, (1997), p.431
9. L. N. Da Costa et al., ApJ 437 L1 (1994)
10. H. Tadros, G. Efstathiou, G. Dalton, MNRAS 296 995 (1998)
11. E. Gaztañaga and C. M. Baugh, MNRAS 294 229 (1998)
12. P. T. P. Viana and A. R. Liddle, MNRAS 281 323 (1996)
13. N. A. Bahcall, X. H. Fan and R. Y. Cen, ApJ 485L 53 (1997)
14. T. Kolatt and A. Dekel, ApJ 479 592 (1997)
15. J. A. Peacock and S. J. Dodds, MNRAS 267 1020 (1994).
16. J. A. Peacock and S. J. Dodds, MNRAS 280 L19 (1996).
17. A. Blanchard, J. Bartlett and R. Sadat preprint astro-ph/9809182 (1998)
18. M. Balogh, A. Babul and D. R. Patton, preprint astro-ph/9809159 (1998)
19. W. Hu, D. Eisenstein, and M. Tegmark, Phys. Rev. Lett., 80 5255 (1998)
20. Y. Fukuda et al., preprint hep-ex/9805021, Phys. Rev. Lett. in press (1998)
21. J. N. Bahcall, P. I. Krastev, and A. Yu. Smirnov, preprint hep-ph/9807216, (1998)
22. C. Athanassopoulos et al., Phys. Rev. Lett. 77 3082 (1996)
23. J. Lesgourgues, D. Polarski and A. A. Starobinsky, MNRAS 297 769 (1998)
24. N. Sugiyama and J. Silk, preprint (1998)




Figure 1: Constraints from LSS and CMB on Λ CDM model




Figure 2: Λ CDM model with $\Omega_\nu = 0.05$. A blue tilt of the primordial power spectrum ($n = 1.3$) is necessary to counteract the damping of small-scale perturbations by free-streaming of the massive neutrinos, which makes the peak of the model fall even farther below that of the data unless $n > 1$. Even with this best-fit value of n , the fit to the data is worse than with no HDM, because CMB observations disfavor such a high value of n .




Figure 3: Λ CDM model with $\Omega_\nu = 0.10$. The best-fit value of n is now 1.5. The fit to the data worsens as more HDM is added.




Figure 4: Constraints from LSS and CMB on ΛCDM model with a broad enhancement centered at $k = 0.06h^{-1}\text{Mpc}$ added to the primordial power spectrum.